



[News](#) [Front Page](#) [Today's Summaries](#) [Latest Headlines](#) [Search Archives](#) [Email Newsletters](#) [RSS Newsfeeds](#) [News Sections](#) [Health & Medicine](#) [Mind & Brain](#) [Plants & Animals](#) [Space & Time](#) [Earth & Climate](#) [Matter & Energy](#) [Computers & Math](#) [Fossils & Ruins](#) [Articles](#) [Videos](#) [Images](#) [Books](#)

Science News

Drug Used To Prevent HIV Transmission From Mother To Child Damages DNA

ScienceDaily (Apr. 8, 2007) — **Studies demonstrate that AZT causes genetic damage that may increase future cancer risk.**

HIV transmission from mother to child can occur in utero, during labor or from breastfeeding. If left untreated, approximately 25 percent of newborns exposed to the virus from their infected mothers will become infected themselves and potentially develop AIDS. Fortunately, antiretroviral drug combinations, which typically include AZT (zidovudine), a nucleoside reverse transcriptase inhibitor (NRTI), have reduced the rate of transmission from mother to child to less than 2 percent in infants who are not breast fed.

NRTIs work by inhibiting the viral reverse transcriptase and by incorporating into the viral DNA and terminating nascent strands, thus preventing the virus from duplicating. However, previous research has shown that NRTIs also incorporate into the DNA of host cells, causing damage that could have long-term health consequences for those exposed to the drugs.

Two new animal studies have examined the cancer-causing effects of transplacental exposure to AZT in mice and rats and found increased rates of tumors and tumors with gene changes that frequently occur in human cancer. In addition, two human studies are the first to observe the induction of mutations and large scale chromosomal damage in red blood cells of newborns exposed to NRTIs in utero.

These, and other, studies were published in April 2007 in a special issue of *Environmental and Molecular Mutagenesis* that examines the latest research on DNA damage and potential health risks related to the use of NRTIs. Besides the effects of NRTIs on nuclear DNA and cancer risk, the issue also contains recent findings on the toxicity of these drugs toward mitochondrial DNA.

Researchers led by Dale M. Walker of Experimental Pathology Laboratories in Herndon, VA, administered AZT in varying doses to female mice and rats during the last 7 days of gestation and examined the tissue of their offspring two years later. They found clear evidence of an AZT-induced increase in the incidence of hemangiosarcoma (cancer originating in cells that line the blood vessels) in male mice and mononuclear cell leukemia in female rats.

There was also some evidence of increased liver cancer and reproductive tumors. "Although the implications of these findings for the long-term health of human children exposed transplacentally to AZT are uncertain, the possibility of increased cancer risk for a subset of these children in mid and late adulthood appears highly plausible," the authors state.

The carcinogenic effects of AZT were further demonstrated by a study on mice led by Hue-Hua

Hong of the National Institute of Environmental Health Sciences in Research Triangle Park, NC. This study found mutations in the K-ras and p53 cancer genes that are often mutated in human lung tumors. The development of lung cancer in these mice suggests that the incorporation of AZT or its metabolites into DNA, oxidative stress, and genomic instability may be the contributing factors to the pattern of mutations observed in the study, according to the authors. They conclude, "The cumulative mutagenesis data suggest that infants exposed transplacentally to AZT may be at increased risk for cancer as they age."

In the first of the two human studies, researchers led by Patricia A. Escobar of the University of Pittsburgh, in Pittsburgh, PA, measured DNA damage caused by AZT in the blood of newborns. They found increased frequencies of glycophorin A mutations in the red blood cells of newborns who had been exposed to AZT plus lamivudine (another type of NRTI) in utero, and these changes persisted for the most part through one year of age. The researchers note that although the combination of the two NRTIs is more effective at preventing transmission of HIV from mother to fetus, it is also more genotoxic than AZT alone. They conclude that "there is a need for careful monitoring of the future health of children who received peripartum AZT-based therapies, the development of new safer NRTIs, and the identification of antimutagenic drugs that will mitigate the side effects of NRTI-based highly active antiretroviral therapy."

In the second study involving humans, researchers led by Kristine L. Witt of the National Institute of Environmental Health Sciences in Research Triangle Park, NC measured the frequency of immature red blood cells (reticulocytes; RET) containing micronuclei (MN), indicators of chromosomal damage, in blood samples of HIV-infected women and their infants exposed to antiretroviral drugs during pregnancy. Most, but not all, of the prenatal treatment regimens in this study included AZT. At birth, the researchers observed ten-fold increases in the frequencies of micronucleated reticulocytes (MN-RET) in the women and infants whose prenatal drug regimen included AZT.

No increases were detected in the women and infants who did not receive prenatal AZT. The frequency of MN-RET in the AZT-exposed newborns decreased over the first 6 months of life to levels seen in nonexposed infants. These findings imply a strong potential for AZT-induced genetic damage in the developing fetus. The authors state "We are concerned about the long-term health implications for these infants because the MN increases noted in this study add to the growing body of evidence that ZDV [AZT] readily induces genetic damage," The authors conclude by emphasizing that they do not advocate eliminating the use of AZT in the treatment of HIV because it is highly effective in preventing mother to child transmission of the virus. However they recommend long-term monitoring of AZT-exposed infants who are HIV uninfected.

Articles:

"Genotoxicity Assessed by the Comet and GPA Assays Following In Vitro Exposure of Human Lymphoblastoid Cells (H9) or Perinatal Exposure of Mother-Child Pairs to AZT or AZT-3TC", Patricia A. Escobar, Ofelia A. Olivero, Nancy A. Wade, Elaine J. Abrams, Carol J. Nesel, Roberta B. Ness, Richard D. Day, Billy W. Day, Quanxin Meng, J. Patrick O'Neill, Dale M. Walker, Miriam C. Poirier, Vernon E. Walker, William L. Bigbee, *Environmental and Molecular Mutagenesis*, **April 2007**, 48:330-343.

"Genetic Alterations in K-ras and p53 Cancer Genes in Lung Neoplasms From Swiss (CD-1) Male Mice Exposed Transplacentally to AZT", Hue-Hua L. Hong, June Dunnick, Ronald Herbert, Theodora R. Devereux, Yongbaek Kim, Robert C. Sills, *Environmental and Molecular Mutagenesis*, **April 2007**, 48:299-306.

"Transplacental Carcinogenicity of 3- β -Azido-3- β -Deoxythymidine in B6C3F1 Mice and F344 Rats", Dale M. Walker, David E. Malarkey, Steven K. Seilkop, Frederick A. Ruecker, Kathleen A. Funk, Marilyn J. Wolfe, Christopher P. Treanor, Julie F. Foley, Fletcher F. Hahn, Jerry F. Hardisty, Vernon E. Walker, *Environmental and Molecular Mutagenesis*, **April 2007**, 48:283-298.

"Elevated Frequencies of Micronucleated Erythrocytes in Infants Exposed to Zidovudine In Utero and Postpartum to Prevent Mother-to-Child Transmission of HIV", Kristine L. Witt, Coleen K. Cunningham, Kristine B. Patterson, Grace E. Kissling, Stephen D. Dertinger, Elizabeth Livingston, Jack B. Bishop, Environmental and Molecular Mutagenesis, **April 2007**, 48:322-329.

See Also:

Health & Medicine

[HIV and AIDS](#)

[Lung Cancer](#)

[Brain Tumor](#)

Plants & Animals

[Mice](#)

[Biotechnology](#)

[Genetics](#)

Reference

[Antiviral drug](#)

[Antiretroviral drug](#)

[Hepatitis B](#)

[Vector \(biology\)](#)

Related Stories

[Zambian Study Finds Longer Breastfeeding Best for HIV-Infected Mothers](#) (Jan. 19, 2010) — A new study from Zambia suggests that halting breastfeeding early causes more harm than good for children not infected with HIV who are born to HIV-positive mothers. Stopping breastfeeding before 18 ... > [read more](#)

[Resistant HIV Quickly Hides In Infants' Cells](#) (May 1, 2007) — New evidence shows that drug-resistant virus passed from mother-to-child can quickly establish itself in infants' CD4+ T cells where it can hide for years, likely limiting their options for future ... > [read more](#)

[Antiretroviral Therapy May Prevent HIV Transmission From Breastfeeding Mothers To Infants](#) (July 31, 2005) — Two new studies support the hypothesis that combination antiretroviral drug therapy may reduce the risk of mother-to-child HIV transmission through breastfeeding, findings that could have significant ... > [read more](#)

[Studies Clarify Risk Factors For Mother-to-Child Transmission Of Hepatitis C Virus](#) (Oct. 31, 2005) — Breastfeeding does not raise the risk of mother-to-child transmission of hepatitis C virus (HCV), according to two new studies published in the December 1 issue of The Journal of Infectious Diseases, ... > [read more](#)

[Maternal HIV-1 Treatment Protects Against Transmission To Newborns](#) (Oct. 19, 2009) — Mothers receiving highly active antiretroviral therapy to treat HIV-1 infection are less likely than untreated mothers to transmit the virus to their newborns through breastfeeding, according to a ... > [read more](#)

[HIV Drug Can Persist In Mothers' Milk, Increasing Risk To Them And Their Babies](#) (Aug. 6, 2008) — A drug commonly used in the developing world to prevent transmission of HIV from mother to child persists in the breast milk and blood of the mothers, putting them and their babies at risk for ... > [read more](#)

[HIV Drug That Protects a Fetus Should Be Avoided for One Year After Childbirth, Researchers Say](#) (Feb. 28, 2010) — A new study found that while nevirapine works well to prevent mother-to-child HIV transmission, a single dose of nevirapine in infected pregnant women can trigger resistance to some forms of the ... > [read more](#)

[Antiretroviral Therapy For Babies, Mothers Safely Reduces HIV Transmission](#) (July 22, 2009) — Daily doses of antiretroviral syrup to breastfeeding infants or treating their HIV-infected mothers with highly active ART drugs is safe and effective in preventing mother-to-child HIV transmission ... > [read more](#)

Just In:

[Mysterious Flash and Missing Belt on Jupiter](#)
[Songbirds Learn Their Songs During Sleep](#)
[Fly Cells Flock Together, Follow the Light](#)
[Ocean Changes May Have Dire Impact on People](#)
[Quantum Dots: Highly Efficient Solar Cells?](#)
[Constraining the Reign of Ancient Egypt](#)
[Astronomers Witness a Star Being Born](#)
[Gulf Oil Spill: Mississippi River May Help](#)

Science Video News



Cleaning Infected Blood

Infectious disease experts designed a machine called the hemopurifier. It works much like a dialysis machine, using thin fibers to capture and remove. ... > [full story](#)

- [Psychobiologists Find Genetic Component in Children's Food Preference](#)
- [Psychologists Test Effects of Household Noise on Children's Verbal Development](#)
- [Pathologists Invent Easy Way To Diagnose MRSA](#)
- [more science videos](#)

Breaking News... from NewsDaily.com

[Yellow sub finds clues to Antarctic glacier's thaw](#)
[Cleared forests lead to rise in malaria in Brazil](#)
[Cord blood transplants a viable option in leukemia](#)
[Americans, Russian blast off for space station](#)
[There's more water on the moon than anyone thought](#)

[more science news](#)

In Other News ...

- [Chinese yuan under scrutiny before G20 meeting](#)
- [Komorowski leads race to Polish presidency](#)
- [BP estimates oil spill up to 100,00 bpd in document](#)
- [Kyrgyz forces remove barricades but Uzbeks still wary](#)
- [Saudi spends \\$1.6 billion to house displaced near Yemen](#)
- [U.N. to cut staff in Afghanistan over security fears](#)
- [New Jersey battles over tax on millionaires](#)
- [Suicide blasts kill 26 at Iraqi state-sector bank](#)
- [more top news](#)